6 - 7 year old check-up



Patient and Family Education

This teaching sheet contains general information only. Talk with your child's doctor or a member of your child's healthcare team about specific care of your child.

TIPS FOR TAKING CARE OF YOUR CHILD

PARENTING

- There are many things you can do as a parent to help your child grow up healthy and well adjusted:
 - Give him lots of love and care.
 - Set fair but firm rules.
 - Enforce the rules each time they are needed.
 - Teach your child to be responsible for his actions and to respect others.
 - Teach him your family values and the difference between right and wrong.
- Your child wants and needs your love and your praise. Hug him, and tell him that you love him often. Tell him when he does something that pleases you. Thank him when he does something nice.
- Your child may want to help around the house. Give him small jobs, such as picking his clothes and toys up off the floor and setting the table.
- Read with your child each day. Let him choose the stories. Visit a nearby library often.
- Doing well in school is important to your child. Talk with your child each day about what happened at school and about his friends. Be positive, and support your child.
 - Make sure he completes any homework.
 - Stay in touch with teachers, and ask for updates on your child's progress.
- Take time out for yourself. Ask someone you trust to watch your child while you do something alone, with your spouse or with a friend.
- Be sure to check out the background of caregivers you do not know.
- If your child does not behave, be calm but firm. Set a good example of how you want him to behave.
 - If he refuses to listen to you, place him in "time out" for 4 to 5 minutes. Have him sit by himself in a safe place. You may also take a privilege away, such as watching a video.
 - Be consistent. Only make rules that you can carry out. If you say you are going to do something, do it.

DEVELOPMENT

- Your child is learning to read and do simple math now. Ask him to read things like signs, ads and
 food wrappers to you. Have him add and subtract spoons, building blocks or other items around the
 house.
- Let your child make some decisions. This makes him feel important and grown-up.
- Make time for your child to have fun. Your child may enjoy doing crafts, sports or learning to swim
 or dance.
- Help your child to follow rules, and expect others to do the same.
- Friends are very important to your child. Provide supervised play times for him with neighbors or school mates.

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.

- Teach your child how to save money for something he wants in the future. Give him a plastic jar, cardboard box or bank to keep his money safe in his room.
- Watch TV with your child and ask him questions about what he sees. Choose shows without violence or sex. Do not allow a TV set in his bedroom.
- Supervise what your child does on the computer. Place the computer in a common area.

SAFETY

- Smoking
 - Do not smoke at home or in the car.
 - Keep your child away from cigarette smoke. Breathing in second-hand smoke can cause asthma, ear infections and other breathing problems.
 - Keep matches and lighters out of reach.
- Home
 - Install smoke and carbon monoxide detectors. Change the batteries twice a year.
 - Teach your child about fire safety. Have a home fire escape plan for your family, and practice it often.
 - Keep cleaning items and medicines locked up and out of reach. Always keep them in their original container.
 - Keep hot items and sharp objects out of reach.
 - Do not let your child play near the stove or use the stove for cooking.
 - Teach him not to open the door for a stranger.
 - Keep guns and bullets locked up and out of reach.

Car

- If your child is less than 4 feet, 9 inches tall, make sure he rides in a booster seat every time you go driving, no matter how short the trip.
- Booster seats make the seat belt fit your child correctly to help prevent injury.
- Follow both the car owner's manual and the child safety seat directions for how to use the seat correctly.
- If your child is more than 4 feet, 9 inches tall, make sure he uses a seatbelt every time you go driving, no matter how short the trip.
- Keep your child in the backseat. Children should ride in the back seat until they are 13 years old.
- Set a good example for your child. Wear your seatbelt, too.

Play

- Spend time playing with your child. Teach him how to share and take turns.
- Help your child use his imagination. Buy toys that help him pretend, think and learn.
- Teach your child safety rules for riding his bike in the park, on the sidewalk and on the street. Make sure your child wears a bike helmet each time he rides.
- Have him wear a helmet and safety gear for other sports, such as skateboarding and roller skating.
- Make sure that your child's play area is safe. Do not allow him to play in the street.
- Teach him how to cross the street safely.
- · Body safety
 - Teach your child:
 - No one should touch his private parts (parts covered by a bathing suit) except to keep him healthy, such as a doctor or nurse during a check-up.
 - Tell a trusted adult if something happens that makes them feel uncomfortable or unhappy.

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.

- Never keep secrets from you unless it is a good surprise, such as a birthday present.
- Animals
 - Teach your child:
 - Ask an adult before petting a strange animal.
 - Do not tease animals or take their food away.
 - Do not put his face near an animal's mouth.
- Outdoors
 - Use a sunscreen and insect repellant that is safe for children. Use it each time he plays or swims outdoors during the day.
 - Make sure your child wears shoes, even in the grass.
 - Teach your child how to swim and be safe in the water. Never leave your child alone in the water. "Swimmies" should not be used as a life vest.

FEEDING

- Eat meals as a family. Turn off the TV and cell phones.
- Keep meal time pleasant. Share news of the day, and talk together. Focus on family news and not on what or how much your child is eating.
- Stick to scheduled meal times and 1 to 2 snacks.
- Your child may enjoy growing a garden, reading about food and helping you cook.
- Your child's doctor may also advise that you give him a multivitamin each day.

Meals

- Your child needs food from all food groups:
 - Lean meat, eggs, beans and cheese for protein
 - Fruits and vegetables for fiber, vitamins and minerals
 - Whole grains, such as cereal, pasta and whole grain bread for fiber, vitamins and minerals. At least half of the grains he eats should be whole grains.
 - Dairy foods, such as milk, cheese and yogurt for calcium.
- Fill half your child's plate with fruits and vegetables.
- Allow your child to choose how much to eat, and do not scold or nag him about what he is eating or not eating.
- Limit fast food meals and snacks. These foods have a lot of fat, sugar, salt and calories but little nutrition.
- If your child takes a lunch to school:
 - Pack him foods from each of the food groups.
 - Let your child help you pack his lunch.
 - Freeze a water bottle, and add it to the lunch box to keep foods cold.

Snacks

- Give your child healthy snacks, such as fruits, raisins, string cheese (cheese sticks), whole wheat crackers, rice cakes and graham crackers.
- Avoid foods with lots of sugar, such as desserts, sweetened cereals and candy. Candy and sticky, sweet foods can cause tooth decay.

Drinks

• Give your child low fat or skim milk.

SLEEP

- Have a regular bedtime and routine for your child.
- Read to your child each night before he goes to sleep.

HEALTHY HABITS

You can help keep your child and your family healthy when you:

- Wash your hands often during the day. Teach your child and other family members to wash their hands often, too. Use warm water and soap and scrub for 20 seconds.
- Have your child bathe regularly. Have him brush his teeth with fluoride toothpaste each morning and at bedtime.
- Take your child to the dentist for regular check-ups.
- Make sure your child gets needed vaccines to help prevent disease.

Your child learns through play. Limit screen time to 1 or 2 hours a day. Some examples of screen time are:

- TV
- Computers
- Videogames
- · Tablet devices
- Smart phones
- Avoid giving fruit punch and 100% fruit juice to your child. It often has extra sugar he does not
 need. Try giving him small pieces of whole fruit instead. If you do give him juice, only give your
 child 4 to 6 ounces a day. Dilute it so it is 1/2 water and 1/2 juice.
- Also avoid giving soda, sports drinks, energy drinks and sweet tea to your child. Give water or diluted juice instead.
- Whether inside or outside, give your child the chance to be active every day. Active play is good for all children. Your child should have 1 hour or more of physical activity daily.

For more information and tips on living a healthy lifestyle, you may go to www.strong4life.com.

FOR YOUR CHILD'S NEXT VISIT

- You and your child's doctor can talk about how things are going with your child.
- Your child will have a physical examination (check-up) that may include getting vaccines. You will
 have a chance to ask questions.
- You may want to talk about your child's:
 - Eating and sleeping
 - Safety
 - Development
 - Behavior, time-out and discipline
 - School, friends and social skills

In case of an urgent concern or emergency, call 911 or go to the nearest emergency department right away.

THINGS FOR YOU TO DO

- Keep a record of when your child is ill or hurt. Share this record with your child's doctors.
- Make a list of questions to talk about with your child's doctor at the next visit.
- Bring your child's shot (vaccine) record to your next visit.



Pediatric Associates, PC Pediatric Associates of Johns Creek, PC

Checkup Schedule

Birth	(In	hosp	ital	H	epatitis	В
-------	-----	------	------	---	----------	---

2-3 Days after Discharge.....Newborn Exam

2 Week.....Exam Only

1 Month......Exam, Hepatitis B

2 Month......Exam, DTaP, Hib, Polio, PCV, Rotavirus

4 Month......Exam, DTaP, Hib, Polio, PCV, Rotavirus

6 Month......Exam, DTaP, Hib, PCV, Rotavirus, CBC (Complete Blood Count)

9 Month......Exam, Hepatitis B, ASQ

12 Month.....Exam, MMR, Varicella, Hepatitis A, CBC

15 Month.....Exam, HIB, PCV

18 Month......Exam, DTaP, Polio, Hepatitis A, CBC, M-CHAT, ASQ

24 Month......Exam, CBC, M-CHAT

30 Month.....Exam, ASQ

3 Year.....Exam, CBC, Urinalysis

4 Year.....Exam, CBC, Urinalysis, Hearing & Vision, DTaP, Polio, MMR, Varicella

5 Year.....Exam, CBC, Urinalysis, Hearing & Vision, DTaP, Polio, MMR, Varicella

(Vaccines given if not completed at 4 year visit.)

6-10 Years....Exam, CBC, Urinalysis, Hearing & Vision as indicated

11 Year.....Exam, CBC, Urinalysis, MCV4, Tdap, HPV*

12-15 Years.. Exam, CBC, Urinalysis, Hearing & Vision as indicated, HPV *

16 Year.....Exam, CBC, Urinalysis, MCV4, Td, Men B, HPV *

17-25 Years...Exam, CBC, Urinalysis, Hearing & Vision as indicated, Men B

- The Ages & Stages Questionnaire (ASQ) screens and assesses the developmental performance of children in the areas of communication, gross
 motor skills, fine motor skills, problem solving, and personal-social skills. It is used to identify children that would benefit from in-depth
 evaluation for developmental delays.
- The M-CHAT is a scientifically validated tool for screening children between 16 and 30 months of age that assesses risk for autism spectrum disorder (ASD).
- A Visual Evoked Potential (VEP) screening is conducted annually on our patients beginning at age six months through 8 years. The purpose of
 this test is the early detection of any abnormalities in your child's vision. We will provide you with further details regarding this non-evasive
 procedure when you arrive for your physical exam. You may also visit www.Diopsys.com to learn more.
- Tdap-(Tetanus, Diphtheria, & Pertussis Vaccine): This vaccine is recommended for children 10 years of age and older. Due to an increase in diagnosed Pertussis (Whooping Cough) cases, this booster is now required for school attendance.
- MCV4-(Meningococcal Vaccine): This vaccine is recommended for children 11 years of age and older. A booster dose is recommended before
 entering college.
- Men B (Meningococcal B)-This vaccine is recommended for patients 16 years of age and older. This vaccine is given in a 2-part series and is
 recommended before entering college.
- We offer the *HPV (Human Papillomavirus vaccine) to both our female and male patients. This vaccine is given in a 2 or 3-part series depending on what age the series is started.

Click here to visit **The Centers For Disease Control and Prevention** website at **www.CDC.gov** to download important vaccine information which may answer any questions you have concerning a particular vaccine.

Dosing On Call

Dosing for infants and children from your healthcare professional

DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.

If possible, use weight to dose; otherwise, use age

Do NOT use with any other product containing acetaminophen.

Infant Active ing	Infants' TYLENOL® Oral Suspension Active ingredient: acetaminophen 160 mg (in each 5 mL)	al Suspension 160 mg (in each 5 mL)	Available in:	Cherry
WEIGHT	6-11 lbs	12-17 lbs	18-23 lbs	
AGE	0-3 months	4-11 months	12-23 months	
	1.25 mL	2.5 mL	3.75 mL	
DOSE	1.25 ml. 2.5 ml. 3.75 ml. 5 ml.	1.25 mt. 2.5 mt. 3.75 mt.	1,25 ml. 3,25 ml. 5 ml.	

Infants' MOTRIN® Concentra' Active ingredient: ibuprofen (NSAID)* 50 mg (ir Nonsteroidal anti-inflammatory drug.	12-17 lbs	6-11 months	1.25 mL	425 mt. 1.25 mt. 1.875 mt.	
Infants' MOTRIN® Active ingredient: ibuprofen (Phonsteroidal anti-inflammatory drug.	WEIGHT	AGE		DOSE	
Consentated Materials Despise	An information from many Lasts up to		Berry Flance	Use product only as directed.	

Available in: We Berry Dye-F	18-23 lbs	12-23 months	1.875 mL	A35 mL 1.255 mL
ited Drops n each 1.25 mL)				

12-17 lbs	18-23 lbs
4-11 months	12-23 months
2.5 mL	3.75 mL
1.25 res. 2.5 res. 5 res.	1.25 mt. 3.25 mt. 5 mt.

2-3 years 24-35 lbs

Dye-Free Cherry

1.25 mi. 2.5 mi. 3.75 mi. 5

5 mL



Children's MOTRIN® (Children's MOTRIN® (Active ingredient: ibuprofen (NSAI Wortin) Weight 24-35 lbs 3 Age 2-3 years 5 mL

9-10 years	6-8 years	4-5 years	2-3 years	AGE
60-71 lbs	48-59 lbs	36-47 lbs	24-35 lbs	WEIGHT

72-95 lbs

Bubble Gurr Dye-Free Berry

ailable in:

11 years 15 mL

15 L



15 mL

12.5 mL

10 mL

7.5 mL 7.5 mL

5 mL 교

DOSE

24-35 lbs

WEIGHT

FILENOL

2-3 years

AGE

15 mL

12.5 mL

10





Children's MOTRIN® Chewables

MOM

Grape 🕝 Bubblegun

Active ingredient: acetaminophen 160 mg (in each chewable tablet)

Children's TYLENOL® Chewables

FILENOL .

Available in:

	"Nonsteroidal anti-ir	WEIGHT 2	AGE 2	ed.
AVAILABLE	hithens		I lon proof set only	as directed

72-95 lbs

60-71 lbs

48-59 lbs

36-47 lbs 4-5 years

24-35 lbs 2-3 years

AGE

11 years

3 tablets

21/2 tablets 9-10 years

2 tablets 6-8 years

11/2 tablets

1 tablet

DOSE

Available in:

Children's TYLENOL Dissolve Packs

Active ingredient: acetaminophen 160 mg (in each pack)

Wild Berry

WEIGHT	24-35 lbs	36-47 lbs	48-59 lbs	9
AGE	2-3 years	4-5 years	6-8 years	9-1
DOSE	1 tablet	1½ tablets	2 tablets	21/2

3 tablets

tablets

72-95 lbs

-71 lbs

11 years

10 years

~
ш
_
m
-
-
Γ
$\overline{}$
11
-
٠.
ò
ပ
ဗ
ဗ
2
S
RUC.
RUC
TRUC.
TRUC
STRUC
STRUC.
ISTRUC.
NSTRUC
NSTRUC.
INSTRUC.
INSTRUC
INSTRUC
T INSTRUC
T INSTRUC
IT INSTRUC
NT INSTRUC
'NT INSTRUC
ANT INSTRUC
ANT INSTRUC
TANT INSTRUC
TANT INSTRUC
TANT INSTRUC
RTANT INSTRUC
RTANT INSTRUC
ORTANT INSTRUC
ORTANT INSTRUC
ORTANT INSTRUC
PORTANT INSTRUC
PORTANT INSTRUC
IPORTANT INSTRUC
MPORTANT INSTRUC
MPORTANT INSTRUC
IMPORTANT INSTRUC

Today's date:

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

- Always read and follow the label on all TYLENOL* and MOTRIN* products.
- · Use only the dosing device (dosing syringe or dosing cup) that came with the product. Do not use any other dosing device.

3 powders

2 powders

2 powders

DOSE

72-95 lbs 11 years

60-71 lbs 9-10 years

48-59 lbs 6-8 years

under 48 lbs under 6 years do not use

PILENOL

NEW!

AGE

All Infants' TYLENOL* and Children's TYLENOL* products have the same strength of acetaminophen: **160 mg** (in each 5 mL, tablet, or pack).

Rely on TYLENOL" and MOTRIN", the two brands of pain and fever relievers that pediatricians recommend most.†

ದ S Alway